

Some Solutions of Berkovits' Superstring Field Theory

J. Klusoň *

*Institute of Theoretical Physics, University of Stockholm, SCFAB
SE- 106 91 Stockholm, Sweden
and
Institutionen för teoretisk fysik
BOX 803, SE- 751 08 Uppsala, Sweden
E-mail: josef.kluson@teorfys.uu.se*

ABSTRACT: In this note we construct exact solution of Berkovits' superstring field theory.

KEYWORDS: String field theory.

*On leave from Masaryk University, Brno

Contents

1. Introduction	1
2. Review of superstring field theory	2
3. General solution in Berkovits string field theory	3
4. Conclusion	8

1. Introduction

Open string field theory is very useful tool for the understanding of the tachyon condensation. (For review and extensive list of references, see [1]). While bosonic string field theory [3] based on the Chern-Simons-like action is very well understood in superstring case the situation is not such a clear. The straightforward superstring extension of the bosonic string field theory [4] suffers from an emergence of contact divergences that forces us to include higher-contact terms when two picture changing operators collide. (For discussion of this issue and other related problems we again recommend to see very good review [1].)

Another interesting approach to the problem of the construction of superstring field theory was given in Berkovits' works [6, 8]. While previous string field theories are based on the Chern-Simons-like actions, Berkovits formulation is based on the Wess-Zummino-Witten-like action. Two major advantages of Berkovits superstring field theory are that it is manifestly $SO(3, 1)$ super-Poincare invariant and that it does not require contact terms to remove tree-level divergences [15] (For recent review of Berkovits superstring field theory (NSFT), see [1, 2, 16].) In particular, it was shown that the calculation of the tachyon potential in BSFT is in a very good agreement with Sen's conjecture [17]. The tachyon kink and lump solution was also analysed (For extensive list of references, see [1, 2].)

In this short note we will construct solution of NSFT following mainly the seminal paper [18]. We begin with the analysis of the equation of motion of the NS sector² of NSFT. This string field theory is characterised with given bulk (bulk on the world-sheet) conformal field theory (CFT) and with given boundary conditions on

²In this paper we will discuss Neveu-Schwarz sector of open string field theory only. However proposal for inclusion of Ramond sector has been done in very interesting paper [7], see also [24].

the boundary of the world-sheet. In other words, in terminology of [13], we have a string field theory characterised by given boundary CFT (BCFT), where string fields belong to the Hilbert space of BCFT. As is well known (for recent discussion of this approach in bosonic open string field theory and extensive list of references, see, for example [10, 12, 13, 22], for case of NSFT see, [20, 21, 24, 25]) fluctuations around field configuration that is solution of the equation of motion are described with the same action as the original one, however with shifted BRST operator. Since this new BRST operator can have very complicated form, there should exist such a field redefinition so that after performing this redefinition in the action, the new BRST operator gains the same structure as the original one, but fluctuation fields are transformed to the fields that belong to the Hilbert space of the new BCFT'. This new BCFT' has the same CFT in the bulk of the world-sheet as BCFT has so that the BRST operator should be the same as the original one, however boundary conditions have changed which is reflected in the change of vertex operators and hence string fields. We will show that the generator of this transformation is in natural way related to the solution of the equation of motion of NSFT so that whenever we can find such a generator, we can very easily find exact solution of the string field equation of motion. We hope that our results could be helpful for finding new exact solutions of the string field theory equation of motion and could be also helpful in present study of the vacuum string field theory [10, 12, 13] (For recent works and extensive list of references, see [26, 27]).

This note is organised as follows. In section (2) we briefly review basic facts about NS sector of Berkovits string field theory.

The main part of this paper is contained in section (3). There we will show that for any solution of the equation of motion of NSFT we can construct new BRST operator according to the analysis [20, 25]. Using particular form of this operator suggested in [19] we will show that there exist field redefinition that maps the new BRST operator to the original one. Under this transformation string fields that took values in Hilbert space of the original BCFT are mapped to the new BCFT' with new boundary conditions. Then we show that the action is invariant under this redefinition. Finally, we extend our analysis to the case of Witten's bosonic open string field theory.

In conclusion (4) we outline our results and suggest possible directions of further research.

2. Review of superstring field theory

In this section we would like to review basic facts about superstring field theory, for more details, see [1, 5, 6, 9]. The general off-shell string field configuration in the GSO(+) NS sector corresponding Grassmann even open string vertex operator Φ of ghost number 0 and picture number 0 in the combined conformal field theory of a

$c = 15$ superconformal matter system, and b, c, β, γ ghost system with $c = -15$. We can also express β, γ in terms of ghost fields ξ, η, ϕ

$$\beta = \partial\xi e^{-\phi}, \quad \gamma = \eta e^{\phi}, \quad (2.1)$$

the ghost number n_g and the picture number n_p assignments are as follows

$$\begin{aligned} b : n_g = -1, n_p = 0 \quad c : n_g = 1, n_p = 0 ; \\ e^{q\phi} : n_g = 0, n_p = q ; \\ \xi : n_q = -1, n_p = 1 \quad \eta : n_q = 1, n_p = -1 . \end{aligned} \quad (2.2)$$

The BRST operator Q_B is given

$$Q_B = \oint dz j(z) = \int dz \left\{ c(T_m + T_{\xi\eta} + T_\phi) + c\partial cb + \eta e^\phi G_m - \eta\partial\eta e^{2\phi} b \right\}, \quad (2.3)$$

where

$$T_{\xi\eta} = \partial\xi\eta, \quad T_\phi = -\frac{1}{2}\partial\phi\partial\phi - \partial^2\phi, \quad (2.4)$$

T_m is a matter stress tensor and G_m is a matter superconformal generator. Throughout this paper we will be working in units $\alpha' = 1$.

The string field action is given [5, 6]

$$S = \frac{1}{2} \int \left((e^{-\Phi} Q_B e^\Phi)(e^{-\Phi} \eta_0 e^\Phi) - \int_0^1 dt (e^{-t\Phi} \partial_t e^{t\Phi}) \left\{ (e^{-t\Phi} Q_B e^{t\Phi}), (e^{-t\Phi} \eta_0 e^{t\Phi}) \right\} \right), \quad (2.5)$$

where $\{A, B\} = AB + BA$ and $e^{-t\Phi} \partial_t e^{t\Phi} = \Phi$. Here the products and integral are defined by Witten's gluing prescription of the string. The exponential of string field is defined in the same manner $e^\Phi = 1 + \Phi + \frac{1}{2}\Phi \star \Phi + \dots$. In the following we will not explicitly write $\star$ symbol. The basis properties of Q_B, η_0 which we will need in our analysis (for more details, see [1] and reference therein) are

$$\begin{aligned} Q_B^2 = 0, \quad \eta_0^2 = 0, \quad \{Q_B, \eta_0\} = 0, \\ Q_B(\Phi_1 \Phi_2) = Q_B(\Phi_1) \Phi_2 + \Phi_1 Q_B(\Phi_2), \\ \eta_0(\Phi_1 \Phi_2) = \eta_0(\Phi_1) \Phi_2 + \Phi_1 \eta_0(\Phi_2), \\ \int Q_B(\dots) = 0, \quad \int \eta_0(\dots) = 0, \end{aligned} \quad (2.6)$$

where Φ_1, Φ_2 are Grassmann even fields.

3. General solution in Berkovits string field theory

Since we will not perform any explicit calculation we will again use abstract Witten's formalism in string field theory [3]. As usual we will not write explicitly the string

field theory star product $\star$. We start with the action

$$S = \frac{1}{2} \int \left((e^{-\Phi} Q_B e^{\Phi}) (e^{-\Phi} \eta_0 e^{\Phi}) - \int_0^1 dt (e^{-t\Phi} \partial_t e^{t\Phi}) \{ (e^{-t\Phi} Q_B e^{t\Phi}), (e^{-t\Phi} \eta_0 e^{t\Phi}) \} \right) . \quad (3.1)$$

We must stress that this action is formulated around one fixed BCFT with given BRST operator Q_B . Let us consider general solution of the equation of motion that arises from the action (3.1)

$$\eta_0(G_0^{-1} Q_B(G_0)) = 0 , \quad G_0 = e^{\Phi_0} . \quad (3.2)$$

Now we would like to study the fluctuation around this solution. For that reason we write general string field containing fluctuation around this solution as

$$G = G_0 h, \quad h = e^{\phi}, \quad G^{-1} = h^{-1} G_0^{-1} . \quad (3.3)$$

To see that this field really describes fluctuations around solution G_0 note that for $\phi = 0, G = G_0$. It is also clear that any string field in the form $e^{\Phi_0 + \phi'}$ can be always rewritten in the form given above.

Inserting this upper expression in (3.1) we obtain an action for ϕ . As was argued in [20] in order to find form of the new BRST operator we must ask a question what form of the equation of motion obeys shifted field $h = e^{\phi}$ ³. Then it was shown [20] that the new BRST operator has a form

$$\tilde{Q}(X) = Q_B(X) + AX - (-1)^X XA , \quad A = G_0^{-1} Q_B(G_0) . \quad (3.4)$$

Let us now presume that the new BRST operator $\tilde{Q}$ can be written as⁴

$$\tilde{Q} = e^{-K} Q_B(e^K) , \quad (3.5)$$

where K is operator of ghost charge and picture number equal to zero that obeys following rules [12]

$$\begin{aligned} K(AB) &= K(A)B + AK(B), \\ \int K(A)B &= - \int AK(B) \end{aligned} \quad (3.6)$$

and commutes with η_0 . In order to have a well defined string field theory, $\tilde{Q}$ must obey (2.6). Firstly, we can easily show that

$$\begin{aligned} \tilde{Q}^2 &= e^{-K} Q_B e^K e^{-K} Q_B e^K = e^{-K} Q_B^2 e^K = 0 , \\ \{\tilde{Q}, \eta_0\} &= \{e^{-K} Q_B e^K, \eta_0\} = e^{-K} \{Q_B, \eta_0\} e^K = 0 , \end{aligned} \quad (3.7)$$

³For alternative and more elegant approach to this problem, see [25].

⁴This suggestion is based on results presented in [19].

since $[K, \eta_0] = 0$. We also have

$$\begin{aligned}
\tilde{Q}(\Phi_1\Phi_2) &= e^{-K} \left[Q_B \left(e^K(\Phi_1\Phi_2) \right) \right] = e^{-K} \left[Q_B \left(e^K(\Phi_1)e^K(\Phi_2) \right) \right] = \\
&= e^{-K} \left[Q_B \left(e^K(\Phi_1) \right) e^K(\Phi_2) + e^K(\Phi_1) Q_B \left(e^K(\Phi_2) \right) \right] = \\
&= e^{-K} \left[Q_B \left(e^K(\Phi_1) \right) \right] e^{-K} e^K(\Phi_2) + e^{-K} e^K(\Phi_1) e^{-K} \left[Q_B \left(e^K(\Phi_2) \right) \right] = \\
&= \tilde{Q}(\Phi_1)\Phi_2 + \Phi_1\tilde{Q}(\Phi_2) ,
\end{aligned} \tag{3.8}$$

where we have used

$$\begin{aligned}
e^K(AB) &= AB + K(A)B + AK(B) + \\
&+ \frac{1}{2}K^2(A)B + K(A)K(B) + \frac{1}{2}AK^2(B) + \dots = e^K(A)e^K(B)
\end{aligned} \tag{3.9}$$

which follows from (3.6). In the same way we can show that

$$e^{-K}(AB) = e^{-K}(A)e^{-K}(B) . \tag{3.10}$$

We can also easily see that (3.5) obeys the last axiom in (2.6)

$$\int \tilde{Q}(X) = \int e^{-K} \left[Q_B \left(e^K(X) \right) \right] \mathcal{I} = \int Q_B \left(e^K(X) \right) e^K(\mathcal{I}) = \int Q_B \left(e^K(X) \right) = 0 , \tag{3.11}$$

where $\mathcal{I}$ is "identity" ghost number and picture number zero field (for recent discussion of some properties of this field, see [11, 27]) which is defined as

$$\mathcal{I}X = X\mathcal{I} = X , \forall X . \tag{3.12}$$

Then we have

$$K(X) = K(\mathcal{I}X) = K(\mathcal{I})X + \mathcal{I}K(X) = K(X) \Rightarrow K(\mathcal{I}) = 0 \tag{3.13}$$

and consequently

$$e^{-K}(\mathcal{I}) = e^K(\mathcal{I}) = \mathcal{I} . \tag{3.14}$$

In (3.11) we have also used

$$\begin{aligned}
\int e^{-K}(X)Y &= \int \sum_n \frac{1}{n!} (-1)^n K^n(X)Y = \\
&= \sum_n - \int \frac{1}{n!} (-1)^n K^{n-1}(X)K(Y) = \sum_n \frac{1}{n!} \int X K^n(Y) = \int X e^K(Y)
\end{aligned} \tag{3.15}$$

which is a result of successive application of the second property of K given in (3.6).

To finish our analysis, we must find such a string field that solves the equation of motion (3.2) and leads to the shifted BRST operator (3.4). In fact, similar calculation has been performed in [21] so that we do not repeat it here. In the same way as in [21] one can show that the field Φ_0 that is a solution of (3.2) and leads to (3.4) has a form

$$\Phi_0 = K^L(\mathcal{I}) , \quad (3.16)$$

where indices R, L corresponds to the left and right side of open string [3, 18].

We have seen that the new operator $\tilde{Q}$ obeys all axioms given in (2.6) and hence we can define NSFT around the new background configuration characterised with this new BRST operator (3.4). It is important to stress that in BCFT language, we have new BRST operator that can be completely general, however the fluctuation fields still belong to the Hilbert space of the original BCFT. For that reason it is natural to find such a field redefinition that maps $\tilde{Q}$ in Q_B so that the new BRST operator is of the same form and the fluctuation fields that belong to the Hilbert space of the original BCFT are mapped to the fields that belong to the Hilbert space of new BCFT'. We will show that this can be easily done in the framework of Berkovits string field theory.

In order to obtain BRST operator of the same functional form as the original one, we perform field redefinition

$$\phi = e^{-K}(\Psi) . \quad (3.17)$$

Using (3.6) we can show

$$\begin{aligned} e^\phi &= \sum_{n=0}^{\infty} \frac{1}{n!} \left(e^{-K}(\Psi) \right)^n = e^{-K} \left(\sum_{n=0}^{\infty} \frac{1}{n!} \Psi^n \right) = e^{-K}(e^\Psi) , \\ e^{-\phi} &= \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} \left(e^{-K}(\Psi) \right)^n = e^{-K} \left(\sum_{n=0}^{\infty} \frac{(-1)^n}{n!} \Psi^n \right) = e^{-K}(e^{-\Psi}) \end{aligned} \quad (3.18)$$

and consequently

$$e^{-K}(e^\Psi) e^{-K}(e^{-\Psi}) = e^{-K}(e^\Psi e^{-\Psi}) = e^{-K}(\mathcal{I}) = \mathcal{I} . \quad (3.19)$$

Using these results we immediately obtain

$$\begin{aligned} & \int (e^{-\phi} \tilde{Q}(e^\phi)) (e^{-\phi} \eta_0(e^\phi)) = \\ &= \int e^{-K}(e^{-\Psi}) e^{-K} Q_B e^K e^{-K}(e^\Psi) e^{-K}(e^{-\Psi}) \eta_0(e^{-K}(e^{-\Psi})) = \\ &= \int e^{-\Psi} e^K \left[e^{-K} Q_B(e^\Psi) e^{-K}(e^{-\Psi}) \eta_0(e^{-K}(e^\Psi)) \right] = \\ &= \int e^{-\Psi} e^K \left[e^{-K}(Q_B(e^\Psi)) e^{-K}(e^{-\Psi} \eta_0(e^\Psi)) \right] = \\ &= \int (e^{-\Psi} Q_B(e^\Psi)) (e^{-\Psi} \eta_0(e^\Psi)) . \end{aligned} \quad (3.20)$$

We can also show that

$$\begin{aligned}
e^{t\phi} &= \sum_n \frac{1}{n!} t^n (e^{-K}(\Psi))^n = e^{-K} \left(\sum_n \frac{1}{n!} (t\Psi)^n \right) = e^{-K}(e^{t\Psi}) , \\
e^{-t\phi} \partial_t e^{t\phi} &= e^{-K} e^{-t\Psi} e^{-K} \partial_t (e^{t\Psi}) = e^{-K} (e^{-t\Psi} \partial_t e^{t\Psi}) , \\
e^{-t\phi} \tilde{Q} e^{t\phi} &= e^{-K} (e^{-t\Psi}) e^{-K} Q_B e^K e^{-K} (e^{t\Psi}) = e^{-K} (e^{-t\Psi} Q_B (e^{t\Psi})) , \\
e^{-t\phi} \eta_0 (e^{t\phi}) &= e^{-K} (e^{-t\Psi} \eta_0 (e^{t\Psi}))
\end{aligned} \tag{3.21}$$

so that we have

$$\begin{aligned}
&\int_0^1 dt (e^{-t\phi} \partial_t e^{t\phi}) \{ (e^{-t\phi} \tilde{Q} (e^{t\phi})), (e^{-t\phi} \eta_0 (e^{t\phi})) \} = \\
&= \int_0^1 dt e^{-K} (e^{-t\Psi} \partial_t e^{t\Psi}) e^{-K} \{ (e^{-t\Psi} Q_B (e^{t\Psi})), (e^{-t\Psi} \eta_0 (e^{t\Psi})) \} = \\
&= \int_0^1 dt (e^{-t\Psi} \partial_t e^{t\Psi}) e^K e^{-K} \{ (e^{-t\Psi} Q_B (e^{t\Psi})), (e^{-t\Psi} \eta_0 (e^{t\Psi})) \} = \\
&= \int_0^1 dt e^{-t\Psi} \partial_t e^{t\Psi} \{ (e^{-t\Psi} Q_B (e^{t\Psi})), (e^{-t\Psi} \eta_0 (e^{t\Psi})) \} .
\end{aligned} \tag{3.22}$$

From (3.20), (3.22) we obtain correct form of the action for fluctuation fields around new background configuration, however with the BRST operator having the same form as the original one. Now the modification of the background is reflected in the change of the string fields that now belong to the Hilbert space of the BCFT' theory.

We can easily previous result to the case of Witten's open bosonic string theory with the action

$$S = \int \frac{1}{2} \Phi Q_B(\Phi) + \frac{1}{3} \Phi \Phi \Phi , \tag{3.23}$$

where now Φ is ghost number one string field. From (3.23) we obtain the equation of motion

$$Q_B \Phi + \Phi \Phi = 0 . \tag{3.24}$$

It is easy to see that the field

$$\Phi_0 = e^{-\Psi_0} Q_B (e^{\Psi_0}) \tag{3.25}$$

with any ghost number zero field Ψ_0 is a solution of the equation of motion. As in previous case we presume that Ψ_0 can be written as follows

$$\Psi_0 = K_L(\mathcal{I}) , \tag{3.26}$$

where K is operator obeying (3.6). Now the new BRST operator has the same form as before

$$e^{-K} Q_B (e^K) . \tag{3.27}$$

Then we can perform the same field transformation as in supersymmetric case and we end up with the action, where the BRST operator has the same form as the original one and the fluctuating field belongs to the Hilbert space of the new BCFT' theory. So that whenever we will find such two BCFT theories that are related field redefinition given above, we can claim that one string field theory action arises from the expansion around the background configuration that is solution of the equation of motion of the original string field theory where the solution can be written as $\Phi_0 = K^L(\mathcal{I})$.

4. Conclusion

In this short note we gave a simple solution of the equation of motion of the Berkovits string field theory. We have shown that whenever we can express shifted BRST operator in the form similar to the form suggested in [19] we can easily obtain the solution of the equation of motion. On the other hand, knowledge of the operator K allows us to find such a solution as well. In particular, operator K can correspond to the generator of broken symmetry by D-brane background as for example, generator of translation in direction transverse to D-brane. Then solution given above corresponds to translation of D-brane in transverse direction. It is clear that these two configurations should be equivalent so that the original and final BCFT should be the same. We will show in forthcoming publication that this is really true.

We have also shown that similar method can be used in case of Witten's bosonic string field theory. However, in this case the appropriate solution corresponds to the pure gauge. The similar result has been discussed recently in [22, 23].

Finally we must stress that our method cannot generate all solutions of equation of motion of string field theory. From discussion given above it is clear that this method is not applicable when we cannot find operator K relating original and final configuration. In particular, this method cannot be used for the case of tachyon condensation.

There are many problems that worth to be studied. Firstly, we will present particular examples, corresponding to the marginal deformation of the original string field theory. Then we will try to formulate our result in the CFT language, using very elegant formalism reviewed recently in [26, 27]. We will return to these problems in future. And finally, we hope to extend our result to the case of NSFT theory including Ramond sector, following recent papers [7, 24].

Acknowledgement

This work is partly supported by EU contract HPRN-CT-2000-00122.

References

- [1] K. Ohmori, "*A Review on Tachyon Condensation in Open String Field Theories*," [hep-th/0102085](#).
- [2] P. J. De Smet, "*Tachyon Condensation: Calculation in String Field Theory*," [hep-th/0109182](#).
- [3] E. Witten, "*Noncommutative Geometry and String Field Theory*," *Nucl. Phys. B* **268** (1986) 253.
- [4] E. Witten, "*Interacting Field Theory of Open Superstrings*," *Nucl. Phys. B* **276** (1986) 291.
- [5] N. Berkovits, "*Super-Poincare Invariant Superstring Field Theory*," *Nucl. Phys. B* **459** (1996) 339, [hep-th/9503099](#).
- [6] N. Berkovits, "*A New Approach to Superstring Field Theory*," [hep-th/9912121](#).
- [7] N. Berkovits, "*The Ramond Sector of Open Superstring Field Theory*," [hep-th/0109100](#).
- [8] N. Berkovits, "*The Tachyon Potential In Open Neveu-Schwarz String Field Theory*," *J. High Energy Phys.* **0004** (2000) 022, [hep-th/0001084](#).
- [9] N. Berkovits, A. Sen and B. Zwiebach, "*Tachyon Condensation in Superstring Field Theory*," *Nucl. Phys. B* **587** (2000) 147, [hep-th/0002211](#).
- [10] L. Rastelli, A. Sen and B. Zwiebach, "*String Field Theory Around the Tachyon Vacuum*," [hep-th/0012251](#).
- [11] I. Ellwood, B. Feng, Y. He and N. Moeller, "*The Identity String Field and the Tachyon Vacuum*," [hep-th/0105024](#).
- [12] L. Rastelli, A. Sen and B. Zwiebach, "*Classical Solutions in String Field Theory Around the Tachyon Vacuum*," [hep-th/0102112](#).
- [13] L. Rastelli, A. Sen and B. Zwiebach, "*Half-strings, Projectors, and Multiple D-branes in Vacuum String Field Theory*," [hep-th/0105058](#).
- [14] L. Rastelli, A. Sen and B. Zwiebach, "*Boundary CFT Construction of D-branes in Vacuum String Field Theory*," [hep-th/0105168](#).
- [15] N. Berkovits and C. T. Echevarria, "*Four-Point Amplitude from Open Superstring Field Theory*," *Phys. Lett. B* **478** (2000) 343, [hep-th/9912120](#).
- [16] N. Berkovits, "*Review of open superstring field theory*," [hep-th/0105230](#).

- [17] A. Sen, "Stable non-BPS bound states of BPS D-branes," *J. High Energy Phys.* **9808** (1998) 010, [hep-th/9805029](#);
"SO(32) spinors of type I and other solitons on brane-antibrane pair," *J. High Energy Phys.* **9809** (1998) 023, [hep-th/9808141](#);
"Type I D-particle and its interactions," *J. High Energy Phys.* **9810** (1998) 021 [hep-th/9809111](#);
"Non-BPS states and branes in string theory," [hep-th/9904207](#), and reference therein.
- [18] G. T. Horowitz, J. Lykken, R. Rohm and A. Strominger, "Purely Cubic Action for String Field Theory," *Phys. Rev. Lett.* **57** (1986) 283.
- [19] J. N. Acosta, N. Berkovits and O. Chandia, "A Note on the Superstring BRST Operator," [hep-th/9902178](#).
- [20] J. Klusoň, "Some Remarks About Berkovits' Superstring Field Theory," [hep-th/0105319](#).
- [21] J. Klusoň, "Proposal for Background Independent Berkovits' Superstring Field Theory," [hep-th/0106107](#).
- [22] T. Takahashi and S. Tanimoto, "Wilson Lines and Classical Solutions in Cubic Open String Field Theory," *Prog. Theor. Phys.* **06** (2001) 863, [hep-th/0107046](#).
- [23] T. Takahashi and S. Tanimoto, "Dilaton Condensation in Cubic Open String Field Theory," [hep-th/0112124](#).
- [24] M. Sakaguchi, "Pregeometrical Formulation of Berkovits' Open RNS Superstring Field Theories," [hep-th/0112135](#).
- [25] M. Marino and R. Schiappa, "Towards Vacuum Superstring Field Theory: The Supersliver," [hep-th/0112231](#).
- [26] D. Gaiotto, L. Rastelli, A. Sen and B. Zwiebach, "Ghost Structure and Closed Strings in Vacuum String Field Theory," [hep-th/0111129](#).
- [27] I. Kishimoto and K. Ohmori, "CFT Description of Identity String Field: Toward Derivation of the VSFT Action," [hep-th/0112169](#).